

Exact finite size spectrum in super sine-Gordon model

Á. Hegedűs^{1,2}, F. Ravanini^{3,1} and J. Suzuki⁴

2nd February 2008

¹*I.N.F.N., Sezione di Bologna - Italy*

²*Research Institute for Particle and Nuclear Physics,
Hungarian Academy of Sciences, Budapest - Hungary*

³*Dip. di Fisica, Università di Bologna - Italy*

⁴*Dept. of Physics, Shizuoka University - Japan*

Abstract

We present the coupled nonlinear integral equations (NLIE) governing the finite size effects in N=1 super sine-Gordon model for the vacuum as well as for the excited states. Their infrared limit correctly yields the scattering data of the super sine-Gordon S-matrix conjectured by Ahn [15]. Ultraviolet analysis is in agreement with the expected conformal data of $c = 3/2$ CFT. Conformal perturbation theory further corroborates this result.

1 Introduction

Finite Size Effects in two dimensional integrable QFT can be studied exactly by means of sets of coupled nonlinear integral equations that can be thought as continuum versions of Bethe Ansatz equations. In particular, the equation introduced for the Sine-Gordon (SG) model by Destri and De Vega [1, 2] through a light-cone lattice regularization and basically equivalent to a similar one found by Klümper, Batchelor and Pearce [3, 4] for the spin 1/2 XXZ chain, once extended to excited states [5, 6], has proven to be a very important tool in establishing and testing the correct connection between the Factorized Scattering description and the Perturbed CFT formulation of SG model [7, 8, 9]. Restricted SG models can be put in correspondence with minimal CFT perturbed by their ϕ_{13} operators in a similar way [10].

It would be a result of principal importance to be able to formulate such non-linear integral equations (NLIE) for many other integrable QFTs. In this paper we report the construction of NLIEs for the N=1 Super Sine-Gordon (SSG) model. The importance of

this generalization lies in the manifest supersymmetry of this integrable model and the link it has with applications in various areas of physics ranging from condensed matter to strings.

To get the NLIEs for SSG model, we proceed by analogy with the pure SG case. There, the NLIE was obtained through a light-cone lattice construction of a 6-vertex model, which is known to be equivalent, from the Bethe Ansatz point of view, to an alternating inhomogeneous spin $1/2$ XXZ chain. In the continuum limit this lattice model is shown to give the equations of motion of SG (or equivalently massive Thirring) model [11, 12]. It is known since a long time that the 6-vertex model on a square lattice, or equivalently spin $1/2$ *homogeneous* XXZ chain, renormalizes on the continuum to a free massless boson, i.e. $c = 1$ CFT [13]. The analog model with spin 1, the integrable spin 1 homogeneous integrable XXZ chain is equivalent to the square lattice 19-vertex model and renormalizes to a $c = 3/2$ boson + Majorana fermion theory which is explicitly $N=1$ supersymmetric.

It is therefore natural to conjecture that the full SSG model can be constructed starting from a light-cone lattice 19-vertex model, or equivalently from a spin 1 alternating inhomogeneous integrable XXZ chain [26]. A direct deduction of the equations of motion, like in [11] or [12] is too cumbersome and beyond the scope of the present paper.

The Bethe Ansatz for spin 1 integrable XXZ chain has been written long time ago [16] and can be generalized with the introduction of an alternating inhomogeneity trivially. In the homogeneous case, a coupled set of two NLIE's equivalent to it on the lattice has been deduced, by use of transfer matrix recurrence relations and T-Q formalism in ref.[14], including excited states. Their generalization to the inhomogeneous case is also trivial. On the other hand a system of two NLIE was conjectured on the continuum by C. Dunning [27] for the vacuum state and checked to give the expected ultraviolet central charge $c = 3/2$. Further analysis of Finite Size Effects in SSG model was carried out in [18].

In section 2 we recall the setup of the model in Lagrangian formulation, as a perturbed CFT and a factorized scattering theory. In section 3 we present the alternating inhomogeneous integrable spin 1 XXZ chain and its Bethe ansatz. The NLIE is presented in section 4 for the spin chain on the lattice and, by performing the scaling limit, in the continuum. Section 5 is devoted to the infrared (IR) analysis reconstructing the supersymmetric solitonic S-matrix. From this section on we limit ourselves mostly to the repulsive regime. The interesting features of the attractive regime, as well as the implications for superminimal model perturbed by Φ_{13} are left for a future investigation. In section 6 the ultraviolet (UV) limit of the NLIE is computed and shown to reproduce data compatible with the expected $c = 3/2$ CFT. A numerical comparison with predictions of conformal perturbation theory is presented in section 7. Finally we draw our conclusions and perspectives for future work in section 8.

2 The super sine-Gordon (SSG) model

The SSG model can be defined on a cylinder of circumference L (thought as spatial direction, with infinite time direction) by the action

$$\mathcal{A}_{SSG} = \int dt \int_0^L dx \int d\vartheta d\bar{\vartheta} (D\Phi \bar{D}\Phi + \mu \cos \beta \Phi)$$

where β is the coupling, μ a bare mass and the superfield Φ expands in components as

$$\Phi(x, t, \theta, \bar{\theta}) = \varphi(x, t) + \vartheta \psi(x, t) + \bar{\vartheta} \bar{\psi}(x, t) + \vartheta \bar{\vartheta} F(x, t)$$

The action in components, once the auxiliary field F is eliminated by the Euler-Lagrange equations of motion, reads

$$\mathcal{A}_{SSG} = \int dt \int_0^L dx \left(\frac{1}{2} \partial_\mu \varphi \partial^\mu \varphi + i \bar{\psi} \gamma^\mu \partial_\mu \psi + \mu \psi \bar{\psi} \cos \frac{\beta}{2} \varphi + \frac{\mu^2}{\beta^2} \cos \beta \varphi \right) \quad (1)$$

This model is known to be integrable since a long time and evidently possesses $N = 1$ supersymmetry. Its ultraviolet (UV) limit is given by a theory of a free massless boson $\Omega(x, t) = \phi(z) + \bar{\phi}(\bar{z}) = \frac{\varphi}{\sqrt{4\pi}}$ (here $z = x + it$, $\bar{z} = x - it$) compactified on a circle of radius $R = \frac{4\sqrt{\pi}}{\beta}$ (i.e. with quasi periodic boundary conditions $\Omega(x + L, t) = \Omega(x, t) + 2\pi m R$ at fixed time, m being called the *winding number*) and a free massless Majorana fermion with antiperiodic (NS = Neveu-Schwarz) or periodic (R = Ramond) boundary conditions $\psi(x + L, t) = \pm \psi(x, t)$. This CFT obviously has central charge $c = \frac{3}{2}$ and a $(U(1) \times \mathbb{Z}_2)_L \times (U(1) \times \mathbb{Z}_2)_R$ symmetry and shows $N = 1$ supersymmetry.

In the NS sector the primary fields are given by vertex operators

$$V_{n,m}^{(r,\bar{r})}(z, \bar{z}) = \bar{\psi}_{\bar{r}} \psi_r : e^{i[(\frac{n}{R} + \frac{mR}{2})\phi(z) + (\frac{n}{R} - \frac{mR}{2})\bar{\phi}(\bar{z})]} :$$

where $r, \bar{r} \in \{0, 1\}$, $\psi_0 = 1$, $\psi_1 = \psi$ of conformal dimensions

$$\Delta_{n,m}^{(r,\bar{r})} = \frac{1}{2} \left(\frac{n}{R} + \frac{mR}{2} \right)^2 + \frac{r}{2} \quad , \quad \bar{\Delta}_{n,m}^{(r,\bar{r})} = \frac{1}{2} \left(\frac{n}{R} - \frac{mR}{2} \right)^2 + \frac{\bar{r}}{2}$$

In the R sector the primary fields are realized by bosonic vertex operators multiplied by the Ising spin field

$$R_{n,m}(z, \bar{z}) = \sigma(z, \bar{z}) : e^{i[(\frac{n}{R} + \frac{mR}{2})\phi(z) + (\frac{n}{R} - \frac{mR}{2})\bar{\phi}(\bar{z})]} :$$

of conformal dimensions

$$\Delta_{n,m} = \frac{1}{2} \left(\frac{n}{R} + \frac{mR}{2} \right)^2 + \frac{1}{16} \quad , \quad \bar{\Delta}_{n,m} = \frac{1}{2} \left(\frac{n}{R} - \frac{mR}{2} \right)^2 + \frac{1}{16}$$

All the other secondary states in the theory can be obtained by applying the Heisenberg algebra and the fermion algebra modes to the primaries. See [18] for details.

The main object encoding the UV operator content of the theory is the modular invariant partition function of the $c = \frac{3}{2}$ CFT, obtained in [17]. It reads

$$Z(R) = \frac{1}{|\eta|^2} \left\{ (\chi_0 \bar{\chi}_{1/2} + \chi_{1/2} \bar{\chi}_0) \sum_{n \in \mathbb{Z} + \frac{1}{2}, m \in 2\mathbb{Z} + 1} + (|\chi_0|^2 + |\chi_{1/2}|^2) \sum_{n \in \mathbb{Z}, m \in 2\mathbb{Z}} + |\chi_{1/16}|^2 \sum_{2n-m \in 2\mathbb{Z} + 1} \right\} q^{\Delta_{n,m}^{(0,0)}} \bar{q}^{\bar{\Delta}_{n,m}^{(0,0)}}$$

where $\eta(q)$ is the Dedekind function and $q = e^{2\pi i \tau}$, τ being the modular parameter. We see from the partition function that in the NS sector the quantum numbers $n, m, r, \bar{r}$ describing the operator content can take the values

$$n \in \mathbb{Z} + \frac{1}{2} \quad m \in 2\mathbb{Z} + 1 \quad \begin{cases} r = 1 & \bar{r} = 0 \\ r = 0 & \bar{r} = 1 \end{cases}$$

or

$$n \in \mathbb{Z} \quad m \in 2\mathbb{Z} \quad \begin{cases} r = \bar{r} = 0 \\ r = \bar{r} = 1 \end{cases}$$

while in the R sector they are constrained to take the values

$$n \in \mathbb{Z} \quad m \in 2\mathbb{Z} + 1$$

or

$$n \in \mathbb{Z} + \frac{1}{2} \quad m \in 2\mathbb{Z}$$

The SSG model can be seen as a perturbation of this $c = 3/2$ CFT by a relevant operator

$$\Phi_{pert} = \sqrt{2}\pi \left(V_{1,0}^{(1,1)} + V_{-1,0}^{(1,1)} \right) = \sqrt{2}\pi : \psi \bar{\psi} \cos \frac{\beta}{2} \varphi : \quad (2)$$

of conformal dimension $\Delta_{pert} = \bar{\Delta}_{pert} = \frac{1}{2} + \frac{\beta^2}{32\pi}$. The other perturbing field

$$\frac{1}{2} \left(V_{(2,0)}^{(0,0)} + V_{(-2,0)}^{(0,0)} \right) =: \cos \beta \varphi :$$

necessary in the classical action to ensure supersymmetry, can be shown to give a contribution to quantum correlators that goes to zero with the cutoff [21] and can be neglected in a quantum formulation. See [18] for a more detailed discussion of these two perturbation terms.

The operator (2) is relevant ($\Delta_{pert} < 1$) if the SSG coupling is in the range $0 < \beta^2 < 16\pi$. At exactly $\beta^2 = 16\pi$ it becomes marginally relevant and describes a perturbation of the WZW $SU(2)_{k=2}$ model by its $J^a \bar{J}^a$ operator.

Opposite to this UV description of SSG as a perturbed CFT, there exists a factorized scattering theory that has been conjectured by C. Ahn [15] to describe the asymptotic

particle properties of SSG model. The S-matrix is written in detail in the original paper [15] and the vacuum structure is carefully analyzed in [18]. While not repeating here those data, we just recall that the S-matrix has a tensor product form

$$S(\theta) = S_{RSOS(2)}(\theta) \otimes S_{SG}(\theta, \beta_{SG})$$

where the RSOS part is the S-matrix of the tricritical Ising model perturbed by ϕ_{13} and describes the supersymmetric structure, while the SG part is the usual solitonic S-matrix of pure sine-Gordon model, at $\beta_{SG}^2 = \frac{16\pi\beta^2}{16\pi+\beta^2}$, describing the bosonic degrees of freedom. The vacuum structure shows, similarly to SG, infinitely many degenerate vacua periodically repeating. The additional RSOS structure amounts to the fact that in each period there are three degenerate vacua, labeled by $\{0, \frac{1}{2}, 1\}$ and connected by the adjacency of the A_3 Dynkin diagram. The elementary excitations are supersymmetric solitons (kinks) that interpolate these vacua [18]. They are the only asymptotic particles of the model if $\frac{16\pi}{3} < \beta^2 \leq 16\pi$ (*repulsive* regime). For $\beta^2 < \frac{16\pi}{3}$ bound states (supersymmetric breathers) appear (*attractive* regime).

3 The inhomogeneous spin 1 XXZ chain and its Bethe ansatz

As said in the introduction, we assume, by analogy with the SG case, that the SSG model can be constructed by taking the appropriate continuum limit of an integrable lattice model: the light-cone lattice 19-vertex model. For a description of this lattice model including its Boltzmann weights, see [17]. Actually, there are infinitely many possible lattice models providing SSG as a continuum limit. Among them, it is convenient for our purposes to choose one that preserves integrability on the lattice, so that the Bethe ansatz machinery can be used to diagonalize its transfer matrix and get the energy levels and in principle other measurable quantities.

The square lattice 19-vertex model, in the limit of continuous time direction while keeping the space direction discretized, is related to the integrable homogeneous spin 1 XXZ chain [24] with anisotropy γ . Its continuum limit is a system of a free massless boson compactified on a radius $R = \sqrt{\pi/(\pi - 2\gamma)}$. The light-cone lattice version of the same model is instead equivalent to a spin 1 XXZ chain, with N sites labeled by $n = 1, \dots, N$, lattice spacing a , anisotropy γ and inhomogeneity $\lambda_n = (-1)^{n+1}\Theta$. The relation between the compactification radius R and the SSG coupling β implies that the anisotropy γ is related to β through $\frac{\gamma}{\pi} = \Delta_{pert}(\beta)$. It is often conveniently parametrized by $\gamma = \frac{\pi}{p+2}$. Throughout this paper, we will mainly focus on the range $\gamma < \frac{\pi}{3}$, i.e. $p > 1$. This corresponds to the repulsive regime of SSG model.

We start from the description of the homogeneous system. At each site n of the chain a 3-dimensional space $V_n \sim \mathbb{C}^3$ is defined, on which the spin operators of the Hamiltonian act. Let $R_{n,m}(\theta)$ be the R matrix acting non trivially on $V_n \otimes V_m$. Its explicit form can be found in [24].

The homogeneous transfer matrix $T_2(\theta)$ is defined by the trace of the product of R over the auxiliary space $V_a \sim \mathbb{C}^3$

$$T_2(\theta) := \text{Tr}_{V_a} R_{a,1}(\theta) \cdots R_{a,N}(\theta).$$

Periodic boundary conditions are imposed here.

The hamiltonian of the associated spin 1 XXZ chain is identified with $\mathcal{H} = \frac{\pi}{\gamma} \frac{d}{d\theta} \log T_2(\theta)$.

$$\mathcal{H} = \sum_{n=1}^N (\sigma_n^\perp - (\sigma_n^\perp)^2 + \cos 2\gamma (\sigma_n^z - (\sigma_n^z)^2) - (2 \cos \gamma - 1)(\sigma_n^\perp \sigma_n^z + \sigma_n^z \sigma_n^\perp) - 4 \sin^2 \gamma (S_n^z)^2) \quad (3)$$

where we write $\sigma_n = S_n \cdot S_{n+1} = \sigma_n^\perp + \sigma_n^z$.

Next we consider the inhomogeneous generalization [1, 2, 26], ($N \in 2\mathbb{Z}$)

$$T_2(\theta|\{\lambda_j\}) := \text{Tr}_{V_a} R_{a,1}(\theta - \lambda_1) \cdots R_{a,N}(\theta - \lambda_N).$$

By following the light-cone approach, we adopt a particular *alternating* choice of the inhomogeneities $\lambda_n = (-1)^n \Theta$ and write $T_2(\theta, \Theta) \equiv T_2(\theta|\{(-1)^n \Theta\})$. The local hamiltonian containing both the two-body and the three-body interaction is then defined by

$$H = \frac{\pi}{\gamma} \frac{d}{d\theta} \log T_2(\theta)|_{\theta=\Theta}$$

The inhomogeneity introduces the interaction between the left-going frame and the right-going one.

Conveniently, we introduce an auxiliary transfer matrix which is defined in a similar manner to T_2 , replacing the spin $1 \times \text{spin } 1$ R by the spin $1 \times \text{spin } \frac{1}{2}$ R matrix. A twist ω at the boundary can also easily be incorporated. Periodic boundary conditions correspond to $\omega = 0$, antiperiodic ones to $\omega = \pi/2$. The explicit eigenvalues of $T_1(\theta)$ and $T_2(\theta)$ then read

$$\begin{aligned} T_1(\theta, \Theta) &= e^{i\omega} \rho(\theta - i\pi) \frac{Q(\theta + i\pi)}{Q(\theta)} + e^{-i\omega} \rho(\theta + i\pi) \frac{Q(\theta - i\pi)}{Q(\theta)} \\ T_2(\theta, \Theta) &= e^{2i\omega} \rho(\theta - i\frac{\pi}{2}) \rho(\theta - i\frac{3\pi}{2}) \frac{Q(\theta + i\frac{3\pi}{2})}{Q(\theta - i\frac{\pi}{2})} + \rho(\theta - i\frac{\pi}{2}) \rho(\theta + i\frac{\pi}{2}) \frac{Q(\theta + i\frac{3\pi}{2}) Q(\theta - i\frac{3\pi}{2})}{Q(\theta + i\frac{\pi}{2}) Q(\theta - i\frac{\pi}{2})} \\ &\quad + e^{-2i\omega} \rho(\theta + i\frac{\pi}{2}) \rho(\theta + i\frac{3\pi}{2}) \frac{Q(\theta - i\frac{3\pi}{2})}{Q(\theta + i\frac{\pi}{2})} \end{aligned}$$

where

$$\rho(z) = \sinh^{N/2} \frac{\gamma}{\pi} (z - \Theta) \sinh^{N/2} \frac{\gamma}{\pi} (z + \Theta)$$

The important function Q is given by the Bethe ansatz roots θ_j , ($j = 1, \dots, M$); $Q(z) = \prod_{j=1}^M \sinh \frac{\gamma}{\pi} (z - \theta_j)$.

Bethe Ansatz for the homogeneous case was written in [16]. For the alternating inhomogeneous case it can be generalized easily, resulting in

$$\left(\frac{\sinh \frac{\gamma}{\pi}(\theta_j + \Theta + i\pi) \sinh \frac{\gamma}{\pi}(\theta_j - \Theta + i\pi)}{\sinh \frac{\gamma}{\pi}(\theta_j + \Theta - i\pi) \sinh \frac{\gamma}{\pi}(\theta_j - \Theta - i\pi)} \right)^{N/2} = -e^{2i\omega} \prod_{k=1}^M \frac{\sinh \frac{\gamma}{\pi}(\theta_j - \theta_k + i\pi)}{\sinh \frac{\gamma}{\pi}(\theta_j - \theta_k - i\pi)}$$

The number of roots M is related to the (conserved) total 3-rd component of the spin S_z of the chain and the number of sites N by

$$S_z = N - M$$

The Bethe equations have a periodicity for $\theta \rightarrow \theta + \frac{i\pi^2}{\gamma} = \theta + i\pi(p+2)$. It is convenient to consider as fundamental strip $-\frac{\pi(p+2)}{2} < \text{Im}\theta \leq \frac{\pi(p+2)}{2}$.

The vacuum state is realized by a maximal set of (quasi)-two-strings (see Fig. 1)

$$\theta_j = \text{Re}\theta_j \pm i \left(\frac{\pi}{2} + \epsilon_j \right)$$

where the deviations $\epsilon_j > 0$ from the pure two-string behavior are assumed to be small and limited by a positive parameter ϵ : $\epsilon_j \leq \epsilon$. The distribution of these roots is denser in the regions near $\pm\Theta$, where also the ϵ_j are minimal. We exemplify this in some plots, at $p = \frac{1}{2}$ (attractive) and $p = \frac{3}{2}$ (repulsive) in appendix C.

The excited states are usually characterized by roots located in positions different than those of the vacuum, as well as some holes. The following classification of the Bethe ansatz roots, other than 2-strings, is possible:

1. inner roots : $|\text{Im}\theta_j| < \frac{\pi}{2}$, $j = 1, \dots, M_I$
2. close roots: $\frac{\pi}{2} < |\text{Im}\theta_j| < \frac{3\pi}{2}$, $j = 1, \dots, M_C$
3. wide roots: $\frac{3\pi}{2} < |\text{Im}\theta_j| < \frac{\pi^2}{2\gamma}$, $j = 1, \dots, M_W$
4. self conjugate roots : $|\text{Im}\theta_j| = \frac{\pi^2}{2\gamma}$, $j = 1, \dots, M_p$.

Besides these roots, the real zeroes of the transfer matrices T_1 and T_2 also play an important role in the description of the excited states. The zeros of transfer matrix $T_2(\theta)$ are called *holes* because they are in 1:1 correspondence with actual holes in the quasi 2-string distribution.

Once the Bethe roots are calculated as solutions to the Bethe equations, the formula

$$e^{i\frac{a}{2}(E \pm P)} = e^{\pm i\omega} \prod_{j=1}^M \frac{\sinh \frac{\gamma}{\pi}(i\pi \pm \theta_j - \Theta)}{\sinh \frac{\gamma}{\pi}(i\pi \mp \theta_j - \Theta)}$$

yields the energy level and total momentum of the corresponding state. Higher integrals of motion can also be expressed in a similar way.

We remark that the generalization of the above formulation to general values of spin is immediate.

4 The NLIE for the inhomogeneous spin 1 XXZ chain

NLIE on the lattice for a homogeneous spin 1 XXZ chain with periodic boundary conditions has been written in [14], to which we refer for notations and presentation of the deduction. The generalization to the alternating inhomogeneous lattice model is quite straightforward and leads, along similar lines, to the following coupled NLIE

$$\begin{aligned}\ln b(\theta) &= C_b + iD_b(\theta + i\epsilon) + ig_1(\theta) + ig_b(\theta) + (G * \ln B)(\theta) - (G^{[2\epsilon]} * \ln \bar{B})(\theta) \\ &\quad + (K^{[-\frac{\pi}{2} + \epsilon]} * \ln Y)(\theta) \\ \ln y(\theta) &= C_y + g_y(\theta) + (K^{[\frac{\pi}{2} - \epsilon]} * \ln B)(\theta) + (K^{[-\frac{\pi}{2} + \epsilon]} * \ln \bar{B})(\theta)\end{aligned}\tag{4}$$

where $B(\theta) = 1 + b(\theta)$, $Y(\theta) = 1 + y(\theta)$ and for any function f and g

$$(f * g)(x) \equiv \int_{-\infty}^{+\infty} dy f(x - y) g(y) \quad , \quad f^{[\pm\eta]}(x) = f(x \pm i\eta)\tag{5}$$

The kernel functions G and K read

$$G(\theta) = \int_{-\infty}^{\infty} \frac{dq}{2\pi} e^{iq\theta} \frac{\sinh \frac{\pi(p-1)q}{2}}{2 \sinh \frac{\pi pq}{2} \cosh \frac{\pi q}{2}} \quad K(\theta) = \frac{1}{2\pi \cosh(\theta)}.$$

We also introduce odd primitives of kernel functions,

$$\chi(\theta) = 2\pi \int_0^\theta dx G(x) \quad \chi_K(\theta) = 2\pi \int_0^\theta dx K(x)$$

that are important in writing the source terms containing information on the excitations:

$$\begin{aligned}g_b(\theta) &= \sum_{j=1}^{N_H} \chi(\theta - h_j) - 2 \sum_{j=1}^{N_H^S} \chi(\theta - h_j^S) - \sum_{j=1}^{M_S} (\chi(\theta - s_j) + \chi(\theta - \bar{s}_j)) \\ &\quad - \sum_{j=1}^{M_C} \chi(\theta - c_j) - \sum_{j=1}^{M_W} \chi_{II}(\theta - w_j) - \sum_{j=1}^{M_{sc}} \chi_{II}(\theta - w_{sc}^{(j)}), \\ g_1(\theta) &= \sum_{j=1}^{N_1} \chi_K(\theta - h_j^{(1)}), \\ g_y(\theta) &= \lim_{\eta \rightarrow 0^+} \tilde{g}_y \left(\theta + i\frac{\pi}{2} - i\eta \right) \\ \tilde{g}_y(\theta) &= i \left\{ \sum_{j=1}^{N_H} \chi_K(\theta - h_j) - 2 \sum_{j=1}^{N_H^S} \chi_K(\theta - h_j^S) - \sum_{j=1}^{M_S} (\chi_K(\theta - s_j) + \chi_K(\theta - \bar{s}_j)) \right. \\ &\quad \left. - \sum_{j=1}^{M_C} \chi_K(\theta - c_j) \right\}\end{aligned}$$

where the second determination function, for $p > 1$

$$\chi_{II}(\theta) = -i \left[\ln \sinh \left(\frac{\theta}{p} \right) - \ln \sinh \left(\frac{\theta + \text{Sign}(\text{Im}\theta)i\pi}{p} \right) \right]$$

is defined by $\chi(x) + \chi(x - i\pi \text{sgn}(x))$.

The driving term bulk contribution exists only in the equation for $\ln b(\theta)$ and it reads

$$D_b(\theta) = N \arctan \frac{\sinh \theta}{\cosh \Theta}$$

The constants C_b and C_y are determined by the asymptotic behavior of both sides of the NLIE

$$C_b = i\hat{C}_b \quad \hat{C}_b = \pi\delta_b + \alpha \quad \alpha = \omega \left(1 + \frac{2}{p} \right) + \chi_\infty (N_- - N_+). \quad (6)$$

$$\delta_b \in \{0, 1\} \quad \text{and} \quad \chi_\infty = \chi(+\infty) = \frac{\pi}{2} \left(1 - \frac{1}{p} \right). \quad (7)$$

$$N_\pm = \left\lfloor 3 \left(\frac{S_z}{p+2} \mp \frac{\omega}{\pi} \right) \right\rfloor - \left\lfloor \frac{S_z}{p+2} \mp \frac{\omega}{\pi} \right\rfloor \quad (8)$$

where $\lfloor x \rfloor$ stands for the integer part of x and

$$C_y = i\pi\delta_y + i\pi(S + M_W + M_{sc}) \quad \delta_y = N_- \bmod 2 \quad (9)$$

We need one more equation for the determination of wide and self-conjugated roots. For the cases $\pi < \text{Im}\theta \leq \frac{\pi}{2}(p+1)$:

$$\ln \tilde{a}(\theta) = C_{\tilde{a}} + ig_{\tilde{a}}(\theta) + (G_{II}^{[-e]} * \ln B)(\theta) - (G_{II}^{[e]} * \ln \bar{B})(\theta), \quad (10)$$

where

$$\begin{aligned} g_{\tilde{a}}(\theta) = & \sum_{j=1}^{N_H} \chi_{II}(\theta - h_j) - 2 \sum_{j=1}^{N_H^S} \chi_{II}(\theta - h_j^S) - \sum_{j=1}^{M_S} (\chi_{II}(\theta - s_j) + \chi_{II}(\theta - \bar{s}_j)) \\ & - \sum_{j=1}^{M_C} \chi_{II}(\theta - c_j) - \sum_{j=1}^{M_W} [\chi_{II}(\theta - w_j)]_{II} - \sum_{j=1}^{M_{sc}} [\chi_{II}(\theta - w_{sc}^{(j)})]_{II}, \end{aligned} \quad (11)$$

$$C_{\tilde{a}} = i \left\{ 2\omega \left(1 + \frac{2}{p} \right) + \frac{\pi}{p} (N_+ - N_-) \right\}. \quad (12)$$

and the “second-second” determination is defined as in [10].

The parameters $h_j, s_j, c_j, \dots$ in the source terms, i.e. the positions of holes, close, wide roots, etc... can be determined recalling that all these objects annulate the functions $B, 1 + \tilde{a}, Y$, etc... This leads to the following *quantization conditions*

- For holes:

$$\frac{1}{i} \ln b(h_j - i\epsilon) = 2\pi I_{h_j} \quad j = 1, \dots, N_H. \quad (13)$$

- A bit formally for special roots:

$$\frac{1}{i} \ln b(s_j - i\epsilon) = 2\pi I_{s_j} \quad j = 1, \dots, M_S. \quad (14)$$

- For close roots (only for the upper part of the close pair):

$$\frac{1}{i} \ln b(c_j^\dagger - i\epsilon) = 2\pi I_{c_j^\dagger} \quad j = 1, \dots, M_C/2. \quad (15)$$

- For wide roots:

$$\frac{1}{i} \ln \tilde{a}(w_j^\dagger) = 2\pi I_{w_j^\dagger} \quad j = 1, \dots, M_W/2. \quad (16)$$

- For self-conjugated roots:

$$\frac{1}{i} \ln \tilde{a}(w_{sc}^{\dagger(j)}) = 2\pi I_{w_{sc}^{\dagger(j)}} \quad j = 1, \dots, M_{sc}. \quad (17)$$

So far we have determined only the upper part of the complex pairs, but the other parts can be determined by simple complex conjugation.

- For zeroes of $T_1(\theta)$:

$$\frac{1}{i} \ln y_1 \left(h_j^{(1)} - i\frac{\pi}{2} \right) = 2\pi I_{h_j^{(1)}} \quad j = 1, \dots, N_1. \quad (18)$$

All the above quantum numbers I_{α_j} 's are half integers. A state is then identified by a choice of the quantum numbers $(I_{h_j}, I_{c_j}, \dots)$. The NLIE itself can impose constraints on the allowed values some of these quantum numbers.

A comment must be done to explain the emergence of *special* objects (holes and/or roots) h_j^S and $s_j, \bar{s}_j$ in the equations above. Normally the function $Z(\theta) = \text{Im} \ln b(\theta)$ is a monotonically increasing function of θ . It may happen, however, that this monotonicity is locally violated. If it happens that a root or hole θ_j has $Z(\theta_j) < 0$, then we say that it is *special*. In this case the term $\ln B = \ln(1 + b)$ inside the convolution terms goes off-branch, so that $\ln B = \log B + 2\pi i$ (where we use the notation $\log$ to indicate the fundamental branch of the logarithm). The 2π jump is responsible of the emergence of the new terms $\chi(\theta - s) + \chi(\theta - \bar{s})$ (or $2\chi(\theta - h^S)$ for holes) in the source part of the equation. Smoothly varying parameters in the NLIE, if an object that was not a special becomes now special, it is compelled to be accompanied by two new emerging holes. So the number $N_{eff} = N_H - 2N_s$ is conserved by smoothly varying parameters.

A careful analysis of the $\theta \rightarrow \pm\infty$ asymptotics of the NLIE shows that the numbers $N_H, M_C, M_W, \dots$ etc... are not arbitrary, but connected to the total 3-rd component of spin S_z through a set of Diophantine equations, the so called *counting equations*

$$N_H - 2N_S = M_C + 2(S_z + M_W + M_{sc}) - N_+ - N_-, \quad (19)$$

$$N_1 - 2N_R^S = S_z + M_W + M_{sc} + M_C^{(2)} - M_R - \left\lfloor \frac{S_z}{p+2} - \frac{\omega}{\pi} + \frac{1}{2} \right\rfloor - \left\lfloor \frac{S_z}{p+2} + \frac{\omega}{\pi} + \frac{1}{2} \right\rfloor. \quad (20)$$

where $M_C^{(2)}$ is the number of those "close roots" whose imaginary part are either in $\{\pi/2, \pi\}$ or $\{-\pi, -\pi/2\}$. There is an additional parity constraint on N_1

$$N_1 + \left\lfloor \frac{S_z}{p+2} + \frac{\omega}{\pi} + \frac{1}{2} \right\rfloor - \left\lfloor \frac{S_z}{p+2} - \frac{\omega}{\pi} + \frac{1}{2} \right\rfloor = \text{even}. \quad (21)$$

We are interested in the *scaling limit* $N \rightarrow \infty$, $a \rightarrow 0$, while the circumference of the cylinder $L = Na$ remains finite. This limit defines the theory on a continuum cylinder of finite spatial size (circumference) L . Bethe equations become infinite in number in this limit. They must be substituted by some sort of "density of roots" method. However the traditional density approach, based on linear integral equations, is valid only in the large L limit and does not take into account finite size effects. The NLIE approach, instead, is able to fully control the physics on the continuum at any size L .

The lattice NLIE (4) has a sensible scaling limit only if we fine tune the inhomogeneity parameter Θ to go to infinity as $\ln N$ in a very precise way

$$\Theta \sim \ln \frac{2N}{\mathcal{M}L}$$

thus introducing a *mass scale* $\mathcal{M}$. The only variation in the NLIE is in the D_b term that becomes simply

$$D_b(\theta) = \mathcal{M}L \sinh \theta$$

The asymptotic particle interpretation of this term will be clear in the IR analysis of next section. We are in presence of a genuine renormalization procedure, leading to a renormalized quantum field theory on the continuum. In the following it is convenient to introduce the dimensionless scale parameter $\ell = \mathcal{M}L$. The UV physics of the model is reproduced for $\ell \rightarrow 0$ (negligible mass scale), while the IR one corresponds to the "large volume" $\ell \rightarrow \infty$ regime.

The counting equations on the continuum are slightly simplified with respect to the lattice. They read

$$N_H - 2N_S = M_C + 2(S + M_W + M_{sc}), \quad (22)$$

$$N_1 - 2N_R^S = S + M_W + M_{sc} + M_C^{(2)} - M_R - \delta_y, \quad (23)$$

where S is now the topological charge of the continuum theory, connected to the lattice spin S_z is:

$$2S = 2S_z - N_+ - N_- . \quad (24)$$

The terms with integer parts on the rhs. of (20) present on the lattice and coming from the value of $\text{Im}B(\pm\infty)$ become zero in the continuum, where $B(\pm\infty) = 1$. The same happens in the additional parity constraint for N_1 that now simply becomes the requirement that it must be even: $N_1 \in 2\mathbb{Z}$.

The NLIE can be solved numerically with an iterative procedure for a given set of quantum numbers I_k . Once the position of the sources and the functions $\ln b$, $\ln y$ have been calculated, the total energy and momentum of a state can be calculated through the formulae

$$\begin{aligned} E = & \mathcal{M} \left(\sum_{j=1}^{N_H} \cosh(h_j) - 2 \sum_{j=1}^{N_H^S} \cosh(h_j^S) - \sum_{j=1}^{M_S} \{ \cosh(s_j) + \cosh(\bar{s}_j) \} \right. \\ & \left. - \sum_{j=1}^{M_C} \cosh(c_j) + \frac{i}{2\pi} \int_{-\infty}^{+\infty} d\theta \sinh(\theta + i\epsilon) \ln B(\theta) - \frac{i}{2\pi} \int_{-\infty}^{+\infty} d\theta \sinh(\theta - i\epsilon) \ln \bar{B}(\theta) \right) \end{aligned} \quad (25)$$

$$\begin{aligned} P = & \mathcal{M} \left(\sum_{j=1}^{N_H} \sinh(h_j) - 2 \sum_{j=1}^{N_H^S} \sinh(h_j^S) - \sum_{j=1}^{M_S} \{ \sinh(s_j) + \sinh(\bar{s}_j) \} \right. \\ & \left. - \sum_{j=1}^{M_C} \sinh(c_j) + \frac{i}{2\pi} \int_{-\infty}^{+\infty} d\theta \cosh(\theta + i\epsilon) \ln B(\theta) - \frac{i}{2\pi} \int_{-\infty}^{+\infty} d\theta \cosh(\theta - i\epsilon) \ln \bar{B}(\theta) \right) \end{aligned} \quad (26)$$

Notice that wide and self-conjugate root contributions drop from this expressions because $\sinh_{II} \theta = \cosh_{II} \theta = 0$ in the repulsive regime $p > 1$ we are considering here.

5 Large scale analysis and Particle Theory

It is interesting to verify which scattering data can be reproduced by the large ℓ limit of this construction. In the $\ell \rightarrow \infty$ limit the convolution integrals involving B and $\bar{B}$ in both equations are exponentially depressed, as can be easily seen by replacing the leading large ℓ contribution that goes as $\ell \sinh \theta$ into the convolution itself. However, this same reasoning cannot be applied to the other convolution integrals involving Y , as the function y has no exponentially leading term. As a result, the $\ell \rightarrow \infty$ asymptotics of the NLIE looks like

$$-i \ln b(\theta) = \hat{C}_b + \ell \sinh \theta + g_1(\theta) + g_b(\theta) - i(K^{[-\frac{\pi}{2}+\epsilon]} * \ln Y)(\theta) + O(e^{-\ell}) \quad (27)$$

$$\ln y(\theta) = C_y + g_y(\theta) + O(e^{-\ell}) \quad (28)$$

The dominating term of the first equation for large ℓ is $\ell \sinh \theta$, thus showing monotonicity of the real part of the function $-i \ln b$, which in turn implies that for large ℓ there are no special objects: $N_s = 0$.

The second equation is now purely algebraic and can be solved for y and the result can be plugged into the first equation. Defining

$$\mathcal{K}(\theta) \equiv (K^{[-\frac{\pi}{2}]} * \ln(1 + e^{C_y + g_y}))(\theta)$$

we have, putting $\omega = 0$ and letting δ_b take both its values 0 or 1

$$-i \ln b(\theta) = \ell \sinh \theta + g_b(\theta) - i \mathcal{K}(\theta) + \delta_b \pi \quad (29)$$

The energy and momentum expressions also simplify in the same way

$$E = \mathcal{M} \sum_{j=1}^{N_H} \cosh(h_j) - \mathcal{M} \sum_{j=1}^{M_C} \cosh(c_j) + O(e^{-\ell}) \quad (30)$$

$$P = \mathcal{M} \sum_{j=1}^{N_H} \sinh(h_j) - \mathcal{M} \sum_{j=1}^{M_C} \sinh(c_j) + O(e^{-\ell}) \quad (31)$$

Let us first examine the case of N_H holes without any complex pair (other than, of course, the quasi 2-strings of the vacuum sea, that do not contribute here, being confined in the dropping convolution integrals). The energy and momentum are given by

$$E = \mathcal{M} \sum_{j=1}^{N_H} \cosh(h_j) \quad , \quad P = \mathcal{M} \sum_{j=1}^{N_H} \sinh(h_j)$$

We can interpret this result by saying that holes represent a system of asymptotic particles of mass $\mathcal{M}$ and rapidity h_j .

The quantization of particle rapidities for large volume in a factorized scattering theory is given by the formula

$$e^{i \mathcal{M} \ell \sinh \theta_k} \Lambda(\theta_k | \{\theta_i\}) = 1. \quad (32)$$

where $\Lambda(\theta | \{\theta_i\})$ is an eigenvalue of the n -particle color transfer matrix defined by the products of two-body S-matrices

$$T(\theta | \{\theta_i\}) = \prod_{i=1}^n S(\theta - \theta_i)$$

A comparison of (32) with the exponentiated form of (29) and recalling the quantization conditions (13,...,18) shows that in general

$$\Lambda(\theta_k | \{\theta_i\}) = e^{i g_b(\theta_k)} e^{\mathcal{K}(\theta_k)} (-1)^{\delta_b}$$

In the case of two particles, Λ are simply the eigenvalues of the two-body S-matrix. Moreover, if the S-matrix has a tensor product form, also its eigenvalues have. The two particle states are realized by choosing $N_H = 2$. The two hole positions are denoted h_1, h_2 . We recall the well known fact that

$$e^{i\chi(\theta)} = S_0(\theta)$$

where $S_0(\theta)$ is the $S_{++}^{++} = S_{--}^{--}$ element of the SG two-body S-matrix. The counting equation (22) gives a few possible choices

- $S = 1$ and $M_C = M_W = M_{SC} = 0$
- $S = 0$ and $M_C = 2, M_W = M_{SC} = 0$
- $S = 0$ and $M_{SC} = 1, M_C = M_W = 0$

Notice that M_W and M_C can only be even, as the complex roots other than self-conjugate always come in pairs. From the other counting equation (23) one can see that for all the 2-hole states $N_1 = 0$.

A simple inspection case by case of the first factor $e^{ig_b(\theta_k)}$ of Λ shows that it reproduces exactly the SG S-matrix and the quantization of close pair imaginary parts studied in [8]. We conclude that

$$\Lambda = \Lambda_{SG} e^{i\mathcal{K}} (-1)^{\delta_b}$$

The Ahn S-matrix will be reproduced if $e^{i\mathcal{K}} (-1)^{\delta_b}$ gives the four eigenvalues of the RSOS(2) S-matrix. As

$$y(\theta) = (-1)^{\delta_y} \tanh\left(\frac{\theta - h_1}{2}\right) \tanh\left(\frac{\theta - h_2}{2}\right)$$

for the calculation of $\mathcal{K}$ we have to distinguish the two cases $\delta_y = 0$ and $\delta_y = 1$.

In the $\delta_y = 0$ case one gets

$$\mathcal{K}(\theta) = i\pi \{Q_1(\theta - h_{12}) - Q_2(\theta - h_1) - Q_2(\theta - h_2)\}, \quad (33)$$

where $h_{12} = \frac{h_1 + h_2}{2}$ and

$$Q_1(\theta) = -\frac{1}{2\pi} \arctan \sinh(\theta) - \frac{i}{2\pi} \ln \cosh(\theta), \quad (34)$$

$$Q_2(\theta) = -\frac{\chi_2(\theta)}{2\pi} - \frac{i}{2\pi} \ln \cosh\left(\frac{\theta}{2}\right). \quad (35)$$

where $\chi_2(\theta)$ is equal to $\chi(\theta)$ with $p = 2$. Important values from the point of view of the quantization equations are

$$\mathcal{K}(h_1) = \mathcal{R}(h_1 - h_2), \quad (36)$$

$$\mathcal{K}(h_2) = \mathcal{R}(h_2 - h_1), \quad (37)$$

where

$$\mathcal{R}(\theta) = \frac{i}{2}\chi_2(\theta) - \frac{i}{2}\psi_0(\theta), \quad (38)$$

$$\psi_0(\theta) = i \ln \frac{\sinh \frac{i\pi+\theta}{4}}{\sinh \frac{i\pi-\theta}{4}}. \quad (39)$$

The following important identity yields

$$\chi_4(\theta) = \frac{1}{2}\chi_2(\theta) + \frac{1}{2}\psi_0(\theta). \quad (40)$$

where $\chi_4(\theta)$ denotes $\chi(\theta)$ calculated at $p = 4$. This gives two of the four eigenvalues of the RSOS(2) S-matrix, namely

$$\pm e^{i\chi_4(\theta)} \frac{\sinh \left(\frac{i\pi+\theta}{4} \right)}{\sinh \left(\frac{i\pi-\theta}{4} \right)} \quad (41)$$

Along similar lines, the case $\delta_y = 1$ gives the other two eigenvalues

$$\pm e^{i\chi_4(\theta)} \sqrt{\frac{\sinh \left(\frac{i\pi+\theta}{4} \right)}{\sinh \left(\frac{i\pi-\theta}{4} \right)}} \quad (42)$$

thus exhausting the RSOS part of the two-body S-matrix. The complete structure of the solitonic S-matrix of Ahn [15] is reproduced.

We observe that the eigenvalues (41,42) of the RSOS part of the color transfer matrix do not contain the $2^{\theta/2\pi i}$ crossing factors of Ahn's S-matrix. This is because the crossing factors correspond to a gauge transformation of the RSOS transfer matrix, under which the eigenvalues obtainable by finite size effects are invariant [33].

The main surprise of this discussion is the fact that to take into account all the 4 RSOS eigenvalues we had to take into account both values $\delta_y = 0, 1$. Now, from the lattice construction, for the states here examined, only $\delta_y = 0$ is allowed. The case where in the second NLIE δ_y is taken to be 1 cannot be deduced from lattice construction. To say that there exists a NLIE where $\delta_y = 1$ is a conjecture, similar in a certain sense to the one leading to the NLIE for odd topological charge sectors in SG [9]. Allowing this extension of the NLIE does not only reproduce the two missing eigenvalues of RSOS part of the S-matrix, but, as we shall see in the following, also the entire R sector of the UV modular invariant partition function. If there is a viable lattice construction for this choice of δ_y or not remains to be clarified.

6 UV limit and kink approximation

Now we examine the ultraviolet limit of the states described by the NLIE. We follow the standard approach described in detail in [1, 20, 4]. The position of the sources for $\ell \rightarrow 0$ can remain finite (central objects), or they can move towards the two infinities as $\pm \ln \left(\frac{2}{\ell} \right)$

(left/right movers). We introduce the finite parts $\theta_j^{\pm,0}$ of their positions θ_j by subtracting the divergent contribution:

$$\{\theta_j\} \rightarrow \left\{ \theta_j^{\pm} \pm \ln \left(\frac{2}{\ell} \right), \theta_j^0 \right\}.$$

We denote the number of right/left moving and central objects by $N_H^{\pm,0}, N_S^{\pm,0}, M_C^{\pm,0}, \dots$ etc. For later convenience we introduce the right/left moving and central spin given by

$$S^{\pm,0} = \frac{1}{2}(N_H^{\pm,0} - 2N_S^{\pm,0} - M_C^{\pm,0} - 2M_W^{\pm,0} - 2M_{sc}^{\pm,0}). \quad (43)$$

In the UV limit the NLIE splits into three separate equations corresponding to the three asymptotic regions. This is why for all the auxiliary functions of the NLIE (4) we define the so called *kink functions* as

$$F_{\pm}(\theta) = \lim_{\ell \rightarrow 0} F \left(\theta \pm \ln \frac{2}{\ell} \right). \quad F \in \{\log b, \log y, \log \tilde{a}, \dots\}. \quad (44)$$

In the UV limit these kink functions satisfy the so called *kink equations* and the energy and momentum can be expressed by them. Performing the above *kink limit* on our NLIE the kink equations take the form

$$\begin{aligned} \log b_{\pm}(\theta) &= i C_{b\pm}^{(i)} \pm i e^{\pm(\theta+i\epsilon)} + i g_{1\pm}^{(0)}(\theta + i\epsilon) + i g_{b\pm}^{(0)}(\theta + i\epsilon) \\ &+ (G * \ln B_{\pm})(\theta) - (G^{+2\epsilon} * \ln \bar{B}_{\pm})(\theta) + (K^{-\frac{\pi}{2}+\epsilon} * \ln Y_{\pm})(\theta), \end{aligned} \quad (45)$$

$$\log y_{\pm} \left(\theta - i\frac{\pi}{2} \right) = \tilde{C}_{y\pm} + \tilde{g}_{y\pm}^{(0)}(\theta) + (K^{-\epsilon} * \ln B_{\pm})(\theta) - (K^{+\epsilon} * \ln \bar{B}_{\pm})(\theta), \quad (46)$$

$$\log y_{\pm}(\theta) = \lim_{\eta \rightarrow \frac{\pi}{2}^-} \log y_{\pm} \left(\theta - i\frac{\pi}{2} + i\eta \right). \quad (47)$$

$$\ln \tilde{a}_{\pm}(\theta) = C_{\tilde{a}} + i C_{\tilde{a}}^{\pm} + i \tilde{g}_{\tilde{a}\pm}^{(0)}(\theta) + (G_{II}^{-\epsilon} * \ln B_{\pm})(\theta) - (G_{II}^{+\epsilon} * \ln \bar{B}_{\pm})(\theta), \quad (48)$$

where in (45) $i \in \{1, 2\}$ depending on the value of the imaginary part of θ . If $0 < \text{Im}\theta < \pi/2$, then $i = 1$, if $\pi/2 < \text{Im}\theta < \pi$, then $i = 2$. The bulky expressions of the constants and the concrete form of the source functions are collected in appendix A. The finite part of the right and left moving objects can be obtained from the kink functions by imposing quantization conditions very similarly to (13,...,18).

After some usual manipulations [1, 20, 4] it turns out that in the UV limit the energy and momentum can be expressed by a sum of dilogarithm functions with the $\theta \rightarrow \pm\infty$ limiting values of the kink functions in their argument. One of these two limiting values is trivial, namely:

$$b_{\pm}(\pm\infty) = \bar{b}_{\pm}(\pm\infty) = 0, \quad (49)$$

$$y_{\pm}(\pm\infty) = (-1)^{\delta_y}. \quad (50)$$

The other limiting values satisfy a nontrivial coupled nonlinear algebraic equations, the so-called plateau equations [1]:

$$\begin{aligned} \log b_{\pm}(\mp\infty) &= i\hat{C}_b \pm 2i\chi_{\infty}(S - 2S^{\pm}) \pm 2\pi i \hat{l}_W^{\pm} \pm i\pi \left(M_{sc} + \frac{N_1}{2} - N_1^{\pm} \right) \\ &+ \frac{\chi_{\infty}}{\pi} \{ \ln B_{\pm}(\mp\infty) - \ln \bar{B}_{\pm}(\mp\infty) \} + \frac{1}{2} \ln Y_{\pm}(\mp\infty), \end{aligned} \quad (51)$$

$$y_{\pm}(\mp\infty) = (-1)^{2S^{\pm} + \delta_y} (B_{\pm}(\mp\infty) \bar{B}_{\pm}(\mp\infty))^{\frac{1}{2}}, \quad (52)$$

where

$$\hat{l}_W^{\pm} = l_W^{\pm} - \frac{M_W^{\pm}}{2} - M_{sc}^{\pm}, \quad (53)$$

irrelevant from the point of view of the exponent of equation (51). Now we have to solve eqs. (51,52) analytically. Motivated by the TBA formulation of the problem, we take the following Ansatz for the solutions of the plateau equations (51,52):

$$b_{\pm}(\mp\infty) = e^{\pm 3i\rho_{\pm}} 2 \cos(\rho_{\pm}), \quad \bar{b}_{\pm}(\mp\infty) = e^{\mp 3i\rho_{\pm}} 2 \cos(\rho_{\pm}), \quad (54)$$

$$B_{\pm}(\mp\infty) = e^{\pm 2i\rho_{\pm}} \frac{\sin(3\rho_{\pm})}{\sin(\rho_{\pm})}, \quad \bar{B}_{\pm}(\mp\infty) = e^{\mp 2i\rho_{\pm}} \frac{\sin(3\rho_{\pm})}{\sin(\rho_{\pm})}, \quad (55)$$

$$y_{\pm}(\mp\infty) = \frac{\sin(3\rho_{\pm})}{\sin(\rho_{\pm})}, \quad Y_{\pm}(\mp\infty) = 4 \cos(\rho_{\pm})^2 > 0. \quad (56)$$

Since we need only $b_{\pm}(\mp\infty)$ and not the (extended) log of it, we can restrict the allowed values of $\rho_{\pm}$ in the $[0, 2\pi]$ interval. The solutions of the plateau equations formally take the form

$$\rho_{\pm} = \pi \left(k_{\rho}^{\pm} \pm \delta_b \pm \frac{\alpha}{\pi} + \Delta_{\rho_{\pm}} \right) - \frac{\pi}{p+2} \left(2k_{\rho}^{\pm} \pm 2\delta_b \pm 2\frac{\alpha}{\pi} + 3\Delta_{\rho_{\pm}} \right), \quad (57)$$

where

$$\Delta_{\rho_{\pm}} = S - 2S^{\pm} - N_{\rho}^{\pm}, \quad N_{\rho}^{\pm} = \left\lfloor 3\frac{\rho_{\pm}}{\pi} \right\rfloor - \left\lfloor \frac{\rho_{\pm}}{\pi} \right\rfloor, \quad n_{\rho}^{\pm} = \left\lfloor 2\frac{\rho_{\pm}}{\pi} \right\rfloor - \left\lfloor \frac{\rho_{\pm}}{\pi} \right\rfloor. \quad (58)$$

and further constraints must be satisfied by the parity of the integers $k_{\rho}^{\pm}$ and $N_{\rho}^{\pm}$:

$$k_{\rho}^{\pm} = 2l_{\rho}^{\pm} + M_{sc} + \frac{N_1}{2} - N_1^{\pm} - n_{\rho}^{\pm}, \quad l_{\rho}^{\pm} \in \mathbb{Z}, \quad (59)$$

$$N_{\rho}^{\pm} \bmod 2 = 2S^{\pm} + \delta_y \bmod 2 \quad (60)$$

Using the dilogarithm sum rule of appendix B. and putting everything together the conformal weights take the form

$$\Delta^{\pm} = \frac{1}{16} \delta_y + \frac{1}{2} \left(\frac{n_{\pm}}{R} \pm \frac{S}{2} R \right)^2 + \tilde{N}_{\pm} + J_{\pm}, \quad (61)$$

where

$$n_{\pm} = - \left(\delta_b \pm k_{\rho}^{\pm} \pm \frac{3}{2} \Delta_{\rho_{\pm}} \right), \quad (62)$$

$$\tilde{N}_{\pm} = \frac{\hat{N}_{\rho}^{\pm} - \delta_y}{8} \mp S^{\pm} n_{\pm} - \frac{3}{2} (S - S^{\pm}) S^{\pm}, \quad \hat{N}_{\rho}^{\pm} = N_{\rho}^{\pm} \bmod 2. \quad (63)$$

$$\begin{aligned} J_{\pm} &= \pm I_{h^{\pm}} \mp 2I_{h_s^{\pm}} \mp 2I_{s^{\pm}} \mp 2I_{c^{\pm}\uparrow} \mp 2I_{w^{\pm}\uparrow} \mp I_{w_{sc}^{\pm}\uparrow} \pm I_{h^{(1)\pm}} \\ &\mp (S^{\pm} + M_W^{\pm} + M_{sc}^{\pm}) \left(\delta_b \pm \frac{N_1}{2} \mp N_1^{\pm} \pm M_{sc} \mp M_{sc}^{\pm} \right) \\ &- (M_W^+ + M_{sc}^+) (S - S^{\pm}) + K_{\pm}. \end{aligned} \quad (64)$$

where K_+ and K_- are given by

$$K_+ = k_W^+ + M_C^{(2)+} (N_1 - N_1^+) - \frac{N_1^+}{2} \left\{ \delta_y + 2(S + M_W + M_{sc}) + M_C^{(2)} - M_C^{(2)+} \right\} \quad (65)$$

$$K_- = k_W^- + \frac{N_1^-}{2} \left(\delta_y - M_C^{(2)} + M_C^{(2)-} \right) \quad (66)$$

where $k_W^{\pm}$ depend only on the actual configuration of wide roots, and they are integers, if $M_{sc}^{\pm}$ are even, and they are half-integers if $M_{sc}^{\pm}$ are odd.

Analyzing the formulae (61-66) of the UV conformal weights, one can see that S can be identified with the winding number m , the parameter $\delta_y \in \{0, 1\}$ of the NLIE distinguishes the NS and R sectors of the theory, and depending on the state under consideration the sum $N_{\pm} + J_{\pm}$ can be either integer or half-integer. Furthermore it can be easily proven that there is a relation between $n_{\pm}$ of (62) and the winding number S , namely in the NS sector

$$n_{\pm} \in \mathbb{Z} \quad \text{if} \quad S \in 2\mathbb{Z}, \quad (67)$$

$$n_{\pm} \in \mathbb{Z} + \frac{1}{2} \quad \text{if} \quad S \in 2\mathbb{Z} + 1, \quad (68)$$

while in the Ramond sector

$$n_{\pm} \in \mathbb{Z} + \frac{1}{2} \quad \text{if} \quad S \in 2\mathbb{Z}, \quad (69)$$

$$n_{\pm} \in \mathbb{Z} \quad \text{if} \quad S \in 2\mathbb{Z} + 1. \quad (70)$$

in full accordance with what described on the UV operator content in section 2. Moreover it can be proven that in the $\delta_y = 1$ case (R sector) the sum $N_{\pm} + J_{\pm}$ is always integer as it must be in the R sector of a $c = \frac{3}{2}$ CFT. Thus, due to the previous remark and the relations (67-70), the conformal weights (61-66) can be interpreted within the framework of $c = \frac{3}{2}$ CFT as conformal weights appearing in the modular invariant partition function of Di Francesco et al. [17].

7 Comparison with PCFT results

In the previous section we saw that the NLIE (4) describes a modular invariant $c = \frac{3}{2}$ CFT in the UV, but we would like to have more evidence that our coupled NLIEs describe the SSG theory. The simplest check is the numerical comparison of NLIE and perturbed conformal field theory for certain states. This comparison is really important in the R sector where the form of the NLIE is a pure conjecture and can not be obtained from the integrable light-cone lattice regularization of the model.

As we already discussed in section 2, the SSG model can be formulated as a perturbed CFT with the action on the cylinder

$$\mathcal{A} = \mathcal{A}_{3/2} + g \int d^2z \Phi_{pert}(z, \bar{z}), \quad (71)$$

where $\mathcal{A}_{3/2}$ is the action of the $c = \frac{3}{2}$ CFT, $g = \frac{\mu}{2\sqrt{2}\pi}$ is the coupling constant with mass dimension $y = 2(1 - 1/R^2)$ with R being the compactification radius of the conformal normalized boson field and the $\Phi_{pert}(z, \bar{z})$ perturbing operator of eq.(2) is normalized by $\langle \Phi_{pert}(z, \bar{z}) \Phi_{pert}(0, 0) \rangle = |z|^{-2(2-y)}$. In the framework of conformal perturbation theory the energy levels can be represented as a sum of integrals over the n-point functions of the unperturbed CFT [22]. In the case of SSG theory there is no bulk energy contribution on account of supersymmetry, and the perturbation series contains only even powers of the coupling g . Using the exact relation between g and the massgap of the model [23]

$$\frac{g\pi}{\sqrt{8}} \gamma \left(\frac{1}{2} - \frac{\beta^2}{32\pi} \right) = \mathcal{M}^{1-\frac{\beta^2}{16\pi}} \left(\frac{\pi}{4} \frac{\beta^2}{16\pi - \beta^2} \right)^{1-\frac{\beta^2}{16\pi}}, \quad \gamma(x) = \frac{\Gamma(x)}{\Gamma(1-x)}, \quad (72)$$

the perturbation series can be expressed as power series of ℓ^{2y} , where $\ell = \mathcal{M}L$ is the dimensionless volume. The first correction to the energy of a state $|a\rangle$ having conformal dimensions $\Delta_a, \bar{\Delta}_a$ can be written as

$$\frac{6L}{\pi} E_a(\ell) = -\left(\frac{3}{2} - 12(\Delta_a + \bar{\Delta}_a)\right) - C_2^{(a)} \ell^{2y} + O(\ell), \quad (73)$$

where the leading order coefficient reads as

$$C_2^{(a)} = \alpha \frac{I^{(a)}}{\pi} \quad (74)$$

where

$$\alpha = \frac{3}{2} \cdot 8^{2/R^2} \frac{1}{\gamma^2 \left(\frac{1}{2} - \frac{1}{2R^2}\right)} \left(\frac{1}{R^2 - 1} \right)^{2-\frac{2}{R^2}}. \quad (75)$$

contains the massgap formula and $I^{(a)}$ is a simple integral expression

$$I^{(a)} = \int \frac{d^2z}{|z|^y} \langle a | \Phi_{pert}(1, 1) \Phi_{pert}(z, \bar{z}) | a \rangle_{\text{conn}}. \quad (76)$$

In [18] the first nontrivial PCFT corrections were calculated for various states in the NS sector and they were used to test the TCSA near UV, and to check the NLIEs proposed to describe the three ground states of the model. Now, having NLIE for excited states as well, it is worthwhile to check it against PCFT predictions. Unfortunately the comparison is quite restricted on both sides. Actually, the applicability of PCFT, mostly in the R sector, is strongly restricted by IR divergencies. Thus, up to two-kink states the number of IR safe states is very small. On the other hand the NLIE can not be solved numerically with present techniques if special objects appear [1]. Thus, the PCFT-NLIE comparison can be done only for a few states in certain restricted regions of the coupling constants.

In the NS sector, in the zero momentum two-kink states space, the energy of the state $|1-\rangle = \frac{1}{\sqrt{2}} \left(V_{1,0}^{(0,0)}(0,0) - V_{-1,0}^{(0,0)}(0,0) \right) |0\rangle$ with conformal weights $\Delta = \bar{\Delta} = \frac{1}{2R^2}$ can be calculated numerically from NLIE in the region $1 < p < 2$. The PCFT correction of the state was calculated in [18] and is given by

$$I^{1-} = \gamma \left(\frac{1}{2} - \frac{1}{2R^2} \right) \left[\gamma \left(-\frac{1}{R^2} \right) \gamma \left(\frac{1}{2} + \frac{3}{2R^2} \right) - \frac{1}{2} \gamma \left(\frac{1}{2} - \frac{1}{2R^2} \right) \gamma \left(\frac{1}{R^2} \right) \right]. \quad (77)$$

In the language of Bethe Ansatz this state corresponds to a $S = 0$ state with 2-holes, one self-conjugated root, and $\delta_b = \delta_y = 0$. We checked numerically our NLIE against PCFT at a lot of values of the compactification radius R , and in every case we experienced very good agreement. To illustrate the agreement between NLIE and PCFT, the numerical comparison of the quantity $\varepsilon(l) = \frac{6L}{\pi} E_a(l) + (\frac{3}{2} - 12(\Delta_a + \bar{\Delta}_a))$ coming from NLIE and PCFT at the specific $R^2 = \frac{7}{3}$ point can be found in table 1. One can see that the two sets of data approach one another as the volume tends to zero.

In the space of zero-momentum two-kink states of the Ramond sector, because of IR divergencies, the only state for which PCFT can be applied is a charged 2-kink state. At UV, this state corresponds to $|S = 1, R\rangle = R_{0,1}(0,0)|0\rangle$ with conformal weights $\Delta = \bar{\Delta} = \frac{1}{16} + \frac{1}{2} \left(\frac{R}{2} \right)^2$, and the first nontrivial correction to this state is given by

$$I^{S=1,R} = -\frac{\pi}{4} \gamma \left(-\frac{1}{2} + \frac{1}{2R^2} \right) \gamma \left(\frac{1}{2} + \frac{1}{2R^2} \right) \gamma \left(-\frac{1}{R^2} \right). \quad (78)$$

In the language of Bethe Ansatz, this state corresponds to a pure two-hole state with $\delta_b = 0, \delta_y = 1$. The numerical comparison of $\varepsilon(\ell)$ coming from NLIE and PCFT at the specific value $R^2 = \frac{5}{3}$ can be found in table 2. The two sets of data converge as the volume tends to zero.

The very good agreement of NLIE and PCFT data give an additional strong support that the NLIE (4) indeed describes the finite size effects of the SSG model.

8 Conclusions and Perspectives

In the present paper we have presented the full excited states NLIE for the N=1 super-sine-Gordon model, deducing it from a lattice NLIE introduced by Suzuki for the spin 1

ℓ	$\varepsilon(\ell)$ (NLIE)	$\varepsilon(\ell)$ (PCFT)	ℓ^{4y}
0.1	0.213251889(1)	0.2135198827	$5 \cdot 10^{-3}$
0.05	0.096639933(5)	0.096695005	10^{-3}
0.01	0.015365337(5)	0.01536672004	$2 \cdot 10^{-5}$
0.005	0.0069587125(5)	0.006959000	$5 \cdot 10^{-6}$
0.001	0.001105913(5)	0.00110592082	10^{-7}
0.0005	0.00500832(5)	0.005008293	$3 \cdot 10^{-8}$
0.0001	0.000079592(5)	0.00007959153	$7 \cdot 10^{-10}$

Table 1: Numerical comparison of PCFT with NLIE for the state $|1-\rangle$ at $R^2 = \frac{7}{3}$.

ℓ	$\varepsilon(\ell)$ (NLIE)	$\varepsilon(\ell)$ (PCFT)	ℓ^{4y}
0.1	0.4411699(1)	0.42645819	0.025
0.05	0.24978747(1)	0.24493591	$8 \cdot 10^{-3}$
0.01	0.06795829(1)	0.067589069	$6 \cdot 10^{-4}$
0.005	0.03894153(2)	0.03881972	$2 \cdot 10^{-4}$
0.001	0.01072140(3)	0.010712145	10^{-5}
0.0005	0.00615559(2)	0.00615251	$5 \cdot 10^{-6}$
0.0001	0.00016979(1)	0.000169776	$4 \cdot 10^{-7}$

Table 2: Numerical comparison of PCFT with NLIE for the state $|S=1, R\rangle$ at $R^2 = 5/3$.

integrable XXZ chain [14]. We have checked the consistency of this procedure by computing some of the infrared and ultraviolet properties known for such quantum field theory. In particular, the S-matrix proposed by C. Ahn [15] has been reproduced, as well as various properties of the $c = \frac{3}{2}$ CFT that is expected to describe the ultraviolet behavior of the model. Much more work has to be done, however. First of all, contrary to the sine-Gordon case, we lack here of a deduction of the equation of motion from the lattice construction. This keeps the whole construction quite conjectural (although on a very solid ground). A neat deduction of equations of motion from the light-cone lattice construction or alternatively from a bosonization of the alternating inhomogeneous XXZ Hamiltonian could be a very important step forward.

We have investigated in detail only the solitonic sector of the model, by focalizing on the repulsive regime where no breathers appear. The structure of the supersymmetric breathers should be investigated for at least two reasons: it completes the analysis of the asymptotic particle spectrum of the theory and gives access to quantum group restrictions describing superconformal minimal models perturbed by Φ_{13} .

The $N=1$ SSG model is just the first in a whole series of Fractional SSG models [25] that can be seen as perturbations of the conformal models introduced in [17]. The full series of coupled NLIE's generalizing those presented here has been proposed, for the vacuum state, by C. Dunning [27]. These models are evidently mapped, by analogy with SG and SSG, into light-cone higher vertex models like the 44-vertex, etc... or equivalently in alternating inhomogeneous XXZ chains of higher and higher spin. It would be interesting to check Dunning's conjecture and extend NLIEs to the structure of excited states. Two applications of this could be of importance: the quantum group restriction leading to $SU(2)$ -coset theories [30] perturbed by $\phi_{1,3}^{(1)}$ and the large spin limit where this series should make contact with the $N=2$ SSG model, whose finite size effects are of principal importance in the formulation of superstrings propagating in pp-wave background [31].

An even more unifying point of view has been proposed in [28], where two massive coupled NLIEs encompass the whole set of $SU(2)$ -cosets, of FSSG models and their parent theory, the so-called SS-model [32].

Finally an investigation of the SSG model with integrable boundaries, along the lines of what done for SG in [29] would be of interest.

We hope to return to these issues in future.

Acknowledgments

First of all we would like to thank Clare Dunning for her interest in this work during its early stages and for many useful discussions. JS would like to thank people at the University of Bologna for their kind hospitality, and members of the University of Wuppertal for discussions. He acknowledges the financial support by the Ministry of Education of Japan, a Grant-in-Aid for Scientific Research 14540376. FR and AH acknowledge the European network EUCLID (HPRN-CT-2002-00325) and the INFN grant TO12 for partial financial support. AH also thanks Hungarian National Science fund OTKA (under T043159

and T049495). FR also acknowledges NATO grant PST.CLG.980424 for partial financial support.

Appendix A

This appendix is devoted to list the constants and source functions appearing in the kink equations (45-48). The list of values of the constants is as follows

$$C_{b\pm}^{(1)} = \hat{C}_b + 2\chi_\infty (S - S^\pm) \pm \pi (M_W - M_W^\pm + M_{sc} - M_{sc}^\pm) + 2\pi l_W^\pm \pm \frac{\pi}{2} (N_1 - N_1^\pm), \quad (79)$$

$$C_{b+}^{(2)} = C_{b+}^{(1)} - 2\pi(N_1 - N_1^+), \quad C_{b-}^{(2)} = C_{b-}^{(1)}, \quad (80)$$

$$\tilde{C}_{y+} = i\pi \left\{ \delta_y + 2(S + M_W + M_{sc}) + M_C^{(2)} - (S^+ + M_W^+ + M_{sc}^+ + M_C^{(2)+}) \right\}, \quad (81)$$

$$\tilde{C}_{y-} = i\pi \left\{ \delta_y - M_C^{(2)} + (S^- + M_W^- + M_{sc}^- + M_C^{(2)-}) \right\}. \quad (82)$$

$$C_{\bar{a}} = i \left\{ 2\omega \left(1 + \frac{2}{p} \right) + \frac{\pi}{p} (N_+ - N_-) \right\}. \quad (83)$$

$$C_{\bar{a}\pm} = \pm 4\chi_\infty (S - S^\pm) \mp 2\pi(S - S^\pm) + 2\pi q_{\bar{a}}^\pm, \quad (84)$$

The numbers $q_{\bar{a}}^\pm$ and $l_W^\pm$ are integers which appear only in the presence of wide roots, and their actual value depends on the relative position of the wide roots. The form of the source functions of (45-48) are given by

$$g_{1\pm,0}^{(0)}(\theta) = \sum_{j=1}^{N_1^{\pm,0}} \chi_K(\theta - h_j^{(1)\pm,0}), \quad (85)$$

$$\begin{aligned} \tilde{g}_{y\pm,0}^{(0)}(\theta) &= i \left\{ \sum_{j=1}^{N_H^{\pm,0}} \chi_K(\theta - h_j^{\pm,0}) - 2 \sum_{j=1}^{N_H^{S\pm,0}} \chi_K(\theta - h_j^{S,\pm,0}) - \sum_{j=1}^{M_S^{\pm,0}} (\chi_K(\theta - s_j^{\pm,0}) + \chi_K(\theta - \bar{s}_j^{\pm,0})) \right. \\ &\quad \left. - \sum_{j=1}^{M_C^{\pm,0}} \chi_K(\theta - c_j^{\pm,0}) \right\}. \end{aligned} \quad (86)$$

$$\begin{aligned} g_{b\pm,0}^{(0)}(\theta) &= \sum_{j=1}^{N_H^{\pm,0}} \chi(\theta - h_j^{\pm,0}) - 2 \sum_{j=1}^{N_H^{S\pm,0}} \chi(\theta - h_j^{S,\pm,0}) - \sum_{j=1}^{M_S^{\pm,0}} (\chi(\theta - s_j^{\pm,0}) + \chi(\theta - \bar{s}_j^{\pm,0})) \\ &\quad - \sum_{j=1}^{M_C^{\pm,0}} \chi(\theta - c_j^{\pm,0}) - \sum_{j=1}^{M_W^{\pm,0}} \chi_{II}(\theta - w_j^{\pm,0}) - \sum_{j=1}^{M_{sc}^{\pm,0}} \chi_{II}(\theta - w_{sc}^{(j)\pm,0}), \end{aligned} \quad (87)$$

$$\begin{aligned}
\tilde{a}_{\pm,0}^{(0)}(\theta) &= \sum_{j=1}^{N_H^{\pm,0}} \chi_{II}(\theta - h_j^{\pm,0}) - 2 \sum_{j=1}^{N_H^{S\pm,0}} \chi_{II}(\theta - h_j^{S\pm,0}) - \sum_{j=1}^{M_S^{\pm,0}} (\chi_{II}(\theta - s_j^{\pm,0}) + \chi_{II}(\theta - \bar{s}_j^{\pm,0})) \\
&- \sum_{j=1}^{M_C^{\pm,0}} \chi_{II}(\theta - c_j^{\pm,0}) - \sum_{j=1}^{M_W^{\pm,0}} [\chi_{II}(\theta - w_j^{\pm,0})]_{II} - \sum_{j=1}^{M_{sc}^{\pm,0}} [\chi_{II}(\theta - w_{sc}^{(j)\pm,0})]_{II}. \quad (88)
\end{aligned}$$

Appendix B

In section 6, in the calculation of the UV conformal weights, the following dilogarithmic sum must be calculated:

$$S_0(\rho) = 2 \left\{ L_+ [e^{3i\rho} 2 \cos(\rho)] + L_+ [e^{-3i\rho} 2 \cos(\rho)] + L_+ \left[\frac{\sin(3\rho)}{\sin(\rho)} \right] \right\}, \quad (89)$$

where

$$L_+(x) = \frac{1}{2} \int_0^x dy \left\{ \frac{\ln(1+y)}{y} - \frac{\ln y}{1+y} \right\}. \quad (90)$$

The function $S_0(\rho)$ has the following simple properties:

$$S_0(\rho) = S_0(\rho + \pi), \quad S_0(\rho) = S_0(-\rho). \quad (91)$$

After some algebra one can prove that

$$S_0(\rho) = \frac{2\pi^2}{3} + \hat{N}_\rho \left\{ 2\pi \left| \rho - \pi \left[\frac{\rho}{\pi} \right] - \frac{\pi}{2} \right| - \pi^2 \right\} - i \hat{N}_\rho \pi \ln(4 \cos^2 \rho), \quad (92)$$

where

$$\hat{N}_\rho = N_\rho \bmod 2 \quad N_\rho = \left[3 \frac{\rho}{\pi} \right] - \left[\frac{\rho}{\pi} \right], \quad (93)$$

and we made the choice of $\ln(-1) = i\pi$.

Appendix C

We solve the Bethe ansatz equation on the lattice directly for $N = 36$, $\gamma = \frac{\pi}{2.5}$ and for various particular $\ell = 2N e^{-\Theta}$ in the ground state. This corresponds to the attractive regime and $\delta_y = 0$. The Bethe ansatz roots and $\Im m(\ln b)$ are plotted below. Here we use the variable $x = \gamma\theta/\pi$.

One sees that the central regime in the graph of $\ln b$ becomes flatter for smaller ℓ . The slope however seems to be always positive.

With increase of N , the Newton method for finding the Bethe ansatz roots becomes quite unstable. We thus adopt a kind of WKB method to locate them. We plot the result for $N = 128$ in Fig. 3. This again support the positivity of the slope qualitatively.

Figure 1: The Bethe ansatz roots (left) and the corresponding $\Im m(\ln b)$ for $N = 36, \gamma = \pi/2.5, \ell = 1$




Figure 2: The Bethe ansatz roots (left) and the corresponding $\Im m(\ln b)$ for $N = 36, \gamma = \pi/2.5, \ell = 0.1$




Figure 3: The Bethe ansatz roots (left) and the corresponding $\Re(\ln b)$ for $N = 128, \gamma = \pi/2.5, \ell = 0.1$




Figure 4: The Bethe ansatz roots (left) and the corresponding $\Im m(\ln b)$ for $N = 38, \gamma = \pi/2.5, \ell = 0.1$





Figure 5: The Bethe ansatz roots (left) and the corresponding $\Im m(\ln b)$ for $N = 38, \gamma = \pi/3.5, \ell = 0.1$



Next we solve the BAE for $N = 38$. This corresponds to the choice $\delta_b = 1$. The plots in Fig 4 show that the slope near the origin is again positive (modulo jumps) for $\gamma = \pi/2.5$ for smaller values of ℓ .

The plots in the repulsive regime ($\gamma = \pi/3.5$) are given in Fig. 5. The right figure shows the negative slope at the origin. (Imagine the “extended Brillouin zone” and line defined by $\Im m x = \pi$.) This indicates that the upper component θ_j of the quasi 2-string centered at the origin has $\frac{d}{d\theta} \Im m \ln b(\theta)|_{\theta=\theta_j} < 0$, i.e. it is a special object. While in the analog attractive case $N_s = 0$ and for the vacuum where there are no holes $N_{eff} = N_H - 2N_s = 0$, here we have now $N_s = 1$. However, the jumps at $x \sim \pm 0.7$ do not correspond to any pair of roots. Therefore they identify two pairs of holes (they are *not* special, as the derivative of $\ln b$ is positive for them). In total we have now $N_H = 2$, but N_{eff} continues to be 0.

References

- [1] C. Destri and H. J. de Vega, Phys. Rev. Lett. **69** (1992) 2313-2317 – hep-th/9203064
- [2] C. Destri and H. J. de Vega, Nucl. Phys. **B438** (1995) 413-454 – hep-th/9407117
- [3] A. Klümper, M.T. Batchelor and P.A. Pearce, J. Phys. **A24** (1991) 3111-3133
- [4] A. Klümper and P. A. Pearce, J. Stat. Phys. **64** (1991) 13-76; Physica **A183** (1992) 304.
- [5] D. Fioravanti, A. Mariottini, E. Quattrini and F. Ravanini, Phys. Lett. **B390** (1997) 243-251 – hep-th/9608091
- [6] C. Destri and H. J. de Vega, Nucl. Phys. **B504** (1997) 621-664 – hep-th/9701107

- [7] G. Feverati, F. Ravanini and G. Takács, Phys. Lett. **B430** (1998) 264-273 – hep-th/9803104
- [8] G. Feverati, F. Ravanini and G. Takács, Nucl. Phys. **B540** (1999) 543-586 – hep-th/9805117
- [9] G. Feverati, F. Ravanini and G. Takács, Phys. Lett. **B444** (1998) 442-450 – hep-th/9807160
- [10] G. Feverati, F. Ravanini and G. Takács, Nucl. Phys. **B570** (2000) 615-643 – hep-th/9909031
- [11] C. Destri and H. de Vega, Nucl. Phys. **B290** (1987) 363
- [12] C. Destri and T. Segalini Nucl. Phys. **B455** (1995) 759 – hep-th/9506120
- [13] M.P.M. den Nijs, Phys. Rev. **B23** (1981) 6111;
A. Luther and I. Peschel, Phys. Rev. **B12** (1975) 3908;
H.J.F. Knops, Ann. Phys. **128** (1980) 448
- [14] J. Suzuki, J.Phys. **A37** (2004) 11957-11970 – hep-th/0410243
- [15] C. Ahn, Nucl. Phys. **B354** (1991) 57-84
- [16] A. N. Kirillov, N. Yu. Reshetikhin, J. Phys. **A20** (1987) 1565;
H. M. Babujian, A. M. Tsvelick, Nucl. Phys. **B265** [FS15] (1986) 24
- [17] P. Di Francesco, H.Saleur and J.-B. Zuber, Nucl. Phys. **B300** [FS22] (1988) 393
- [18] Z. Bajnok, C. Dunning, L. Palla, G. Takacs and F. Wagner Nucl. Phys. **B679** (2004) 521.
- [19] S. Ferrara, L. Girardello and S. Sciuto, Phys. Lett. **B76** (1978) 303
- [20] Al. B. Zamolodchikov, Nucl. Phys. **B342** (1990) 695;
T. R. Klassen, E. Melzer, Nucl. Phys. **B338** (1990) 485.
- [21] H.C. Liao and P. Mansfield, Nucl. Phys. **B344** (1990) 696.
- [22] T. R. Klassen, E. Melzer, Nucl. Phys. **B350** (1991) 635.
- [23] P. Baseilhac and V.A. Fateev, Nucl. Phys. **B532** (1998) 567 – hep-th/9906010
- [24] V.A. Fateev and A.B. Zamolodchikov, Sov. J. Nucl. Phys. **32** (1981) 298
- [25] D. Bernard and A. LeClair, Phys. Lett. **B247** (1990) 309
- [26] N. Yu. Reshetikhin, H. Saleur Nucl. Phys. **B419** (1994) 507 – hep-th/9309135

- [27] C. Dunning, J. Phys. **A36** (2003) 5463 – hep-th/0210225
- [28] Á. Hegedűs, Nucl.Phys. **B732** (2005) 463-486 – hep-th/0507132
- [29] C. Ahn, M. Bellacosa and F. Ravanini, Phys.Lett. **B595** (2004) 537-546 – hep-th/0312176;
 C. Ahn and R.I. Nepomechie, Nucl.Phys. **B676** (2004) 637-658 – hep-th/0309261
 C. Ahn, Z. Bajnok, R.I. Nepomechie, L. Palla, G. Takacs, Nucl.Phys. **B714** (2005) 307-335 – hep-th/0501047
- [30] F. Ravanini, Mod. Phys. Lett. **3A** (1988) 397
- [31] J. Maldacena and L. Maoz, JHEP **0212** (2002) 046 – hep-th/0207284
- [32] V.A. Fateev, Nucl. Phys. **B473 [FS]** (1996) 509
- [33] T.R. Klassen and E. Melzer, Nucl. Phys. **B382** (1992) 441-485 – hep-th/920203;
 G. Takacs and G. Watts, Nucl. Phys. **B642** (2002) 456-482 – hep-th/020307